



E999

JACC March 12, 2013

Volume 61, Issue 10



## Imaging

### INTRATHORACIC FAT VOLUME IS AN INDEPENDENT PREDICTOR OF CORONARY ARTERY DISEASE IN HEALTHY, ASYMPTOMATIC SUBJECTS

Poster Contributions

Poster Sessions, Expo North

Sunday, March 10, 2013, 9:45 a.m.-10:30 a.m.

Session Title: Imaging: CT/Multimodality VI

Abstract Category: 20. Imaging: CT/Multimodality

Presentation Number: 1230-370

Authors: *Harshal R. Patil, Rajiv Chhabra, James H. O'Keefe, John H. Helzberg, Kevin F. Kennedy, Samantha King, Damini Dey, Randall C. Thompson, Saint Luke's Hospital of Kansas City, Kansas City, MO, USA, University of Missouri Kansas City, Kansas City, MO, USA*

**Background:** Intrathoracic adipose tissue (IAT) appears to be associated with cardiac events in patients, but it is not clear if it is predictive of CAD in asymptomatic subjects.

**Methods:** 348 asymptomatic patients undergoing screening non-contrast CT to determine coronary calcium score (CCS) were studied. IAT volumes were quantified using QFAT software with user-defined range of CT slices covering the heart. We tested 2 nested models. We first created an APRIORI logistic regression model to predict CCS>100 Agatston Units based on traditional risk factors: age, male, smoking, LDL, hypertension, DM, and metabolic syndrome. The next model then added IAT >250 cm<sup>3</sup>. We computed odds ratios for IAT, and compared c-statistics, and computed the integrated discrimination improvement (IDI) for the nested models.

**Results:** Of the 348 subjects, 61 (17.5%) had CCS>100 with an average score of 109.5. When IAT >250 was added to the traditional model, it was independently associated with greater risk of CCS>100 (Odds Ratio 2.07, 95% CI (1.01, 4.23) p=0.046) (Figure 1). Compared to the traditional model, the addition of IAT non-significantly increased the c-statistic (0.771 vs 0.778, p=0.54), However there was a significant value of IDI (Absolute IDI=0.017, Relative IDI=14%, p=0.045). IAT was independently associated with metabolic syndrome (OR: 8.92 95% CI (3.2, 24.9) p<.0001)

**Conclusions:** IAT is an independent predictor of CCS > 100 in asymptomatic subjects and correlates more significantly than traditional risk factors.

